

# Oceanic residual topography agrees with mantle flow predictions at long wavelengths

Ting Yang<sup>1</sup> Louis Moresi<sup>1</sup> R. Dietmar Müller<sup>2</sup> Michael Gurnis<sup>3</sup>

<sup>1</sup>*School of Earth Sciences, University of Melbourne, Melbourne, Vic 3010, Australia*

<sup>2</sup>*Earthbyte Group, School of Geosciences, University of Sydney, Sydney, NSW 2006, Australia*

<sup>3</sup>*Seismological Laboratory, California Institute of Technology, Pasadena, CA 91125, USA*

## Key Points

Synthetic analysis suggests that the long-wavelength residual topography can be recovered from recent point-wise inferences in the ocean.

Appropriate damping factors must be used to recover the long-wavelength residual topography correctly.

Amplitudes (about +/-1km) and patterns of the recovered long-wavelength residual topography from recent oceanic estimates agree with mantle flow predictions.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/2017GL074800

## Abstract

Dynamic topography, the surface deflection induced by sub-lithospheric mantle flow, is an important prediction made by geodynamic models, but there is an apparent disparity between geodynamic model predictions and estimates of residual topography (total topography minus lithospheric and crustal contributions). We generate synthetic global topography fields with different power spectral slopes and spatial patterns to investigate how well the long-wavelength (spherical degree 1 to 3) components can be recovered from a discrete set of samples where residual topography has been recently estimated. Analysis of synthetic topography, along with observed geoid and gravity anomalies, demonstrates the reliability of signal recovery. Appropriate damping factors, which depend on the maximum degree in the spherical harmonic expansion that is used to fit the samples, must be applied to recover the long-wavelength topography correctly; large damping factors smooth the model excessively and suppress residual topography amplitude and power spectra unrealistically. Recovered long-wavelength residual topographies based on recent oceanic point-wise estimates with different spherical expansion degrees agree with each other and with the predicted dynamic topography from mantle flow models. The peak amplitude of the long-wavelength residual topography from oceanic observations is about 1 km, suggesting an important influence of large-scale deep mantle flow.

## Introduction

Mantle convection deflects the lithosphere and generates topography at the earth's surface which is generally referred to as *dynamic topography* [Pekeris, 1935; Parsons and Daly, 1983]. Dynamic topography has an important influence on the gravitational field [Hager and Richards, 1989] and on long-term regional and global sea-level trends [Gurnis, 1993; Lithgow-Bertelloni and Gurnis, 1997; Moucha et al., 2008; Spasojevic and Gurnis, 2012]. It thus forms an important component of reconstructing continental surface environments through time and of understanding the driving forces behind associated transgressions and regressions of inland seas.

There are generally two means of estimating dynamic topography. One is to subtract contributions from the density heterogeneities within the crust and lithosphere (isostatic topography) from the total topography [Cochran and Talwani, 1977; Crough, 1978; Schroeder, 1984; Colin and Fleitout, 1990; Winterbourne et al., 2014]. The derived dynamic topography using this approach is usually referred to as *residual topography*. Alternatively, surface deflections induced by mantle convection can be computed directly by solving the equations for viscous flow. The estimated large-scale dynamic topography based on these two strategies should be approximately consistent if the density and viscosity structures within the lithosphere and mantle are well constrained [Zhang et al., 2012; Yang et al., 2016]. However, because isostatic topography from crustal and lithospheric loads dominates the total topography, and because the density and viscosity structures within the lithosphere and mantle are not well constrained [Le Stunff and Ricard, 1995; Panasyuk and Hager,

2000b], the estimated residual topography and calculated dynamic topography often do not agree. Although mantle flow models often predict large-amplitude negative dynamic topography in east and southeast Asia [Ricard *et al.*, 1993; Lithgow-Bertelloni and Gurnis, 1997; Lithgow - Bertelloni and Richards, 1998; Steinberger, 2007; Zhang *et al.*, 2012; Flament *et al.*, 2013; Yang and Gurnis, 2016; Zhirovic *et al.*, 2016], residual topography has been estimated to be close to zero or positive in the same region [Le Stunff and Ricard, 1995; Panasyuk and Hager, 2000b; Wheeler and White, 2000; Kaban *et al.*, 2004; Flament *et al.*, 2013]. The amplitude of the estimated long-wavelength residual topography generally ranges between 0 and 500 m [Le Stunff and Ricard, 1995; Panasyuk and Hager, 2000b; Wheeler and White, 2000; Kaban *et al.*, 2004], although some have estimated the values to be higher, up to about 1 km [Davies and Pribac, 1993; Gurnis *et al.*, 2000]. In contrast, mantle flow models, which reproduce the observed geoid or stratigraphic records of stable continental interiors, usually predict long-wavelength dynamic topography with an amplitude of about 1 to 2 km [Ricard *et al.*, 1993; Lithgow-Bertelloni and Gurnis, 1997; Lithgow - Bertelloni and Richards, 1998; Steinberger, 2007; Conrad and Husson, 2009; Zhang *et al.*, 2012; Flament *et al.*, 2013; Yang and Gurnis, 2016].

Efforts have been made to reduce predicted topographic amplitudes from mantle flow models so as to fit both the estimated residual topography and geoid observations [Panasyuk and Hager, 2000a; Čadek and Fleitout, 2003; Steinberger, 2007]. Panasyuk and Hager [2000a] suggest that inclusion of one or more low viscosity layers within the mantle transition zone, the amplitude of dynamic topography can be significantly reduced through strongly decoupling lithospheric deflections from deep mantle flow. Although they inferred

radial mantle viscosity profiles by a joint fit of the geoid and residual topography and the fit to the geoid is good, the fit to the estimated residual topography is not, with a variance reduction on residual topography less than 7 % [Panasyuk and Hager, 2000a]. Čadek and Fleitout [2003] inverted for lateral viscosity variations within the upper 300 km of the mantle and predicted a small amplitude dynamic topography with the calculated geoid fitting the observed. However, the pattern of the calculated dynamic topography by Čadek and Fleitout [2003] appears at odds with residual topography estimations. For example, their predicted highest topography is centered on Taiwan in east Asia, inconsistent with residual topography estimates [Panasyuk and Hager, 2000b; Kaban et al., 2004; Steinberger, 2007; Hoggard et al., 2017].

Dynamic topography is especially difficult to estimate on the continents, due to their structural complexity. In contrast, oceanic regions are tectonically simpler and thus have the potential of yielding more precise estimates of residual topography. Recent estimates of residual topography in the oceanic realm [Czarnota et al., 2013; Winterbourne et al., 2014; Hoggard et al., 2016; Hoggard et al., 2017] considered precise contributions from within the crust and lithosphere, including sedimentary layer thickness and density, crustal thickness, and empirical bathymetry-age relationship. These point-wise estimates of residual topography, although sparse and unevenly distributed, has a large-scale pattern and amplitude consistent with the predicted dynamic topography from mantle flow models [Yang and Gurnis, 2016]. However, the extent to which these unevenly-distributed point-wise measurements constrain the pattern and amplitude of long-wavelength residual topography remains a topic of discussion [Hoggard et al., 2017]. Here we address this issue

using a series of synthetic residual topography fields and demonstrate that, with the published observation locations, the long-wavelength (degree 1 to 3) residual topography can be well constrained.

## Method

To investigate the ability of recovering the long-wavelength (degree 1 to 3) residual topography based on observations at sparse, irregularly-distributed points, we first generate random synthetic topography fields and estimate their long-wavelength components by fitting a specific set of discrete samplings with spherical harmonic expansions.

A synthetic topography is generated from a combination of spherical harmonics up to degree 40 (about 1000 km wavelength). At wavelengths less than 1000 km, regional isostasy due to elastic flexure gradually dominates the topography. The synthetic topography  $h$ , is represented as:

$$h(\theta, \varphi) = \sum_{l=1}^{40} \sum_{m=0}^l (C_{lm} \cos \varphi + S_{lm} \sin \varphi) P_l^m(\cos \theta) \quad (1)$$

Here  $\theta$ ,  $\varphi$  are colatitude and longitude, respectively.  $P_l^m$  is a Schmidt semi-normalized associated Legendre polynomial of degree  $l$  and order  $m$ . The spherical harmonic coefficients  $C_{lm}$  and  $S_{lm}$  are generated following uniform random distribution firstly, but

then the amplitude of each degree  $A(l) = \sqrt{\sum_{m=0}^l (C_{lm}^2 + S_{lm}^2)}$  is forced to vary as

$$A(l) = \begin{cases} 300 & l = 1 \\ 1000 \times l^p & l > 1 \end{cases} \quad (2)$$

Both recent estimates of residual topography [Hoggard et al., 2016; Hoggard et al., 2017] and dynamic topography predictions [Yang and Gurnis, 2016] suggest that the amplitude of dynamic topography at degree 1 is much smaller than that at degree 2. The exponent  $p$  controls the slope of the power spectra. By increasing the exponent  $p$ , short-wavelength contributions increase.

We sample the synthetic topography at 2297 points fixed at locations of the Hoggard et al. [2017] data set, with the sampled values deviated from a given synthetic topography by a Gaussian distributed random error. The standard error is 200 m, similar to the observation error estimated by Hoggard et al. [2017], but see more discussions on the measurement error in the supplementary material. Among the 2297 points, 1161 points are better constrained [Hoggard et al., 2017] and thus the Gaussian distributed measurement error is halved. We then carry out spherical harmonic expansions based on these point-wise synthetic topography observations with the least-square method [Hoggard et al., 2016; Yang and Gurnis, 2016]. To suppress unrealistic short-wavelength perturbations and to avoid over fitting the data, some form of damping is often used (for a general description of such problems in geophysics see [Hartzell and Heaton, 1983; Rawlinson and Sambridge, 2003; Tarantola, 2005]). A non-negative damping factor  $\lambda_2$  which mainly suppresses the spatial gradient of the function is employed [Hartzell and Heaton, 1983] in most of our synthetic tests – see supplements for additional discussion. The long-wavelength components can be well recovered by choosing appropriate  $\lambda_2$ .

The power spectrum of the recovered spherical-harmonic expansion is compared against the prescribed synthetic topography power spectrum. The power spectrum at each

degree is defined as the square of the amplitude:

$$P(l) = \sum_{m=0}^l (C_{lm}^2 + S_{lm}^2) \quad (3)$$

We define a variance difference to depict the power spectra difference between the synthetic and recovered topographies at long wavelengths (degree 1 to 3). The variance difference is defined as

$$V_{1-3} = \frac{\sum_{l=1}^3 \sum_{m=0}^l [(C_{lm}^R - C_{lm}^S)^2 + (S_{lm}^R - S_{lm}^S)^2]}{\sum_{l=1}^3 \sum_{m=0}^l [(C_{lm}^S)^2 + (S_{lm}^S)^2]} \quad (4)$$

Where  $S$  and  $R$  represent coefficients of the synthetic and recovered topographies, respectively. Long-wavelength components are well recovered for a small variance difference.

## Results

We start with the long-wavelength (degree 1-3) components recovery from a specific set of point-wise measurements of the synthetic topography. The synthetic topography has an exponent of  $p=-0.3$  which generates substantial short-wavelength perturbations (Fig. 1a) compared to long wavelengths (Fig. 1b). For the same synthetic topography field, we conduct spherical harmonic expansions with different maximum expansion degrees and damping factors  $\lambda_2$ , while evaluating the error of the fit (Fig. 1c-f). The error of the fit to observations decreases with increasing maximum expansion degrees or number of inversion parameters (Fig. 1c). However, beyond degrees about 5, the improvement in the misfit decreases slowly, especially when we consider the number of free parameters. In contrast to



the error of the fit to observations, the average global difference between synthetic and recovered topographies *does not decrease* with increasing maximum expansion degree after 5 (Fig. 1d). This is also true for the global long-wavelength topography difference (Fig. 1e) and variance difference (Fig. 1f). That the global synthetic-recovered topography differences do not decrease with increasing degrees beyond degree 5 indicates that the topography recoverability and resolution is constrained by the number and distribution of observations, and given the present topography observations, increasing the maximum spherical degree from degree 5 (36 parameters) to degree 30 (961 parameters) does not help resolving the global long-wavelength or short-wavelength features. Significantly, the error of the fit to observations and global synthetic-recovered topography differences are sensitive to the damping factor, when we truncate the expansion at a degree larger than about 5. With increasing number of degrees in the expansion, the appropriate damping factor decreases, suggesting that if more parameters are sought from the inversion, then one should not smooth the model excessively.

We fix the maximum expansion degrees to 5, 15, and 30, respectively and investigate the influence of the damping factor on the error of the fit to observations and variance difference (Fig. 2a, b, c) in more detail. The model roughness, defined as the norm of the recovered topography gradient, is large when the model has large-amplitude short-wavelength perturbations. The roughness increases with the decrease of the damping factor. A smooth function has a large error of fit to observations, whereas a rough function usually yields a small error. A suitable compromise is often achieved at the elbow of the misfit-roughness plot [Hartzell and Heaton, 1983; Deschamps et al., 2001; Yue et al., 2013].

With a maximum expansion degree 5 and the damping factor 0.84 that is chosen at the elbow of the misfit-roughness relationship, the predicted long-wavelength topography (Fig. 2a) is comparable with the synthetic one (Fig. 1b), in both pattern and amplitude. A slight change of the damping factor does not change the global topography differences (Fig. 1d, e) or variance difference (Fig. 1f, 2a) significantly. Good estimates of the long-wavelength topography are also found by fixing the maximum expansion degrees to 15 and 30, respectively, but with smaller damping factors (Fig. 2b, c). Fixing the damping factor at 0.84 (elbow of the misfit-roughness relationship in Fig. 2a), the recovered power spectra with different maximum expansions are compared against the synthesis (Fig. 2d). Although the recovered power spectra are close to the actual values when the maximum expansion degree is around 5, they are strongly underestimated when the expansion grows to degree 30, with the peak-to-peak amplitude significantly suppressed (Fig. 2e).

We generate additional random synthetic topography fields which also have the exponent  $p=-0.3$ , but differ significantly in spatial patterns with each other (Fig. S1). The long-wavelength components of these randomly generated synthetic topographies, which have significantly different spatial patterns, can all be recovered, suggesting that the distribution of observations relative to the spatial pattern of topography has only limited influences on the long-wavelength components recovery.

Because the amplitude spectrum of the Earth's dynamic topography is unknown, we evaluate the robustness of synthetic topography recovery in terms of different amplitude spectra. Increasing the exponent  $p$  from  $-0.3$  (Fig. 1) to  $0.0$  significantly increases the short-wavelength synthetic topography contributions (Fig. 3a) while reducing the exponent  $p$

from -0.3 to -0.8 significantly reduces the short-wavelength contributions (Fig. 3f). Although increasing the maximum spherical degree beyond degree about 5 reduces the error of the fit to observations, it does not lead to a better recovery of the long-wavelength synthetic topography (Fig. 3c, h). With appropriate damping factors chosen at the elbow of the misfit-roughness plot (Fig. 3d, i), the recovered long-wavelength topographies (Fig. 3e, j) are consistent with those of the synthetic topographies (Fig. 3b, g).

Although increasing the maximum expansion degree after degree about 5 does not help reducing global synthetic-recovered topography differences, the magnitude of the power spectrum is largely reproduced when appropriate damping factors are chosen (Fig. S2, S3). However, the uncertainty is large and increases with increasing maximum expansion degrees. When large damping factors are used, the magnitude of the recovered power spectra is unrealistically suppressed (Fig. S2). The spectral magnitude is strongly overestimated at short wavelengths when the short-wavelength energy is small (Fig. S3f), especially for large measurement errors, possibly due to the over-fitting of the observations.

We explore the recovery of long-wavelength geoid and free-air gravity anomalies given the specific set of discrete observations. This is a helpful set of experiments as the geoid, gravity anomalies and dynamic topography are closely related [Hager *et al.*, 1984; Yang and Gurnis, 2016] and synthetic topography generally has a different long-wavelength phase and power spectrum from either the geoid or the dynamic topography. The geoid is dominated by long-wavelength components (Fig. S4a, b), while the free-air gravity anomaly has substantial short-wavelength contributions (Fig. S4e, f). Our tests suggest that we are able to recover the long-wavelength components of these fields (Fig. S4d, h), strengthening the

argument that long-wavelength residual topography can be recovered from the sparse estimates by *Hoggard et al.* [2017].

We recover the long-wavelength dynamic topography based on recent point-wise residual topography estimates in the oceanic realm [*Hoggard et al.*, 2017] and compare them with mantle flow predictions [*Yang and Gurnis*, 2016]. Residual topography estimates show a good correlation with the predicted dynamic topography from mantle flow models (Fig. 4a, b). For example, mantle flow models predict large-scale uplift in the central Pacific and around southern Africa, matching large positive residual topography. Mantle flow models predict large depressions around subducted slabs, where the residual topography is also low. We carry out spherical harmonic expansions of the 2297 residual topography observations from *Hoggard et al.* [2017] with different maximum expansion degrees and damping factors (Fig. 4c). With the maximum spherical degree as 5, 12 and 30, we set the corresponding damping factors to 0.55, 0.057 and 0.01, respectively, based on the misfit-roughness relationship (Fig. S5). The estimated long-wavelength (degree 1 to 3) residual topography with different expansion degrees yield similar spatial patterns and amplitudes (Fig. 4d-f) and agrees with mantle flow model predictions (Fig. 4b). The estimated long-wavelength residual topography has a peak-to-peak amplitude of 1 km (Fig. 4d-f), contrary to the results of [*Hoggard et al.*, 2017], who used a combination of gradient regularization damping factor  $\lambda_2$  and amplitude damping factor  $\lambda_1$  which vary between 0.32 and 3.2 and between 100 and 1000, respectively. Using these damping factors, amplitudes of the recovered dynamic topography are significantly smaller (Fig. S6). The estimated residual topography power spectrum is compared with that of dynamic

topography from mantle flow predictions (Fig. S7). Due to the jagged perturbations of the recovered power spectrum around true values (Fig. S2, S3), only first-order features of the recovered power spectra are described. At degrees lower than  $\sim 15$ , spectral magnitudes of the estimated residual topography and dynamic topography are generally consistent. However, at degrees larger than  $\sim 15$ , residual topography spectral magnitude is significantly larger than that of dynamic topography. This inconsistency at short wavelengths may originate from the limitation of mantle flow model predictions (see discussions in *Yang and Gurnis* [2016]) or from limitations in residual topography spectral estimates (Fig. S3f). The residual topography spectral power of degrees larger than 15 is around one order smaller than that of degree 2-3, suggesting a factor of 2 to 3 in amplitude reduction.

## Discussion and Conclusion

Dynamic topography is an important prediction of mantle flow models. However, its amplitude and pattern is controversial. We conduct synthetic experiments and demonstrate that long-wavelength dynamic topography components can be recovered from recent point-wise residual topography observations in the oceanic realm [*Hoggard et al.*, 2017]. Analysis of the geoid and free-air gravity fields further consolidates this argument. Spherical expansions of recent residual topography observations with different maximum degrees and respective appropriate damping factors all yield large-amplitude long-wavelength residual topography that agrees with the mantle flow model predictions.

The prominent uplift around the central Pacific and southwest Indian Ocean and the

depression around southeast Asia that have long been predicted by mantle flow models [Hager and Richards, 1989; Gurnis, 1993; Ricard et al., 1993; Lithgow-Bertelloni and Gurnis, 1997; Lithgow - Bertelloni and Richards, 1998; Steinberger, 2007; Conrad and Husson, 2009; Spasojevic and Gurnis, 2012; Zhang et al., 2012; Flament et al., 2013; Yang and Gurnis, 2016] are well revealed by the recovered long-wavelength residual topography (Fig. 4). Many residual topography models, although with large uncertainties, suggest a large-scale negative dynamic topography around South America [Panasyuk and Hager, 2000b; Kaban et al., 2004; Steinberger, 2007]. Point-wise residual topography observations also show a deep Argentine Basin (Fig. 4a) that is consistent with mantle flow model predictions [Shephard et al., 2012]. However, the amplitude of the negative long-wavelength dynamic topography around South America is small (Fig. 4b), and shallow density heterogeneities may exert additional contributions regionally [Rodríguez Tribaldos et al., 2017], making the recovery of long-wavelength negative dynamic topography around South America more difficult.

Dynamic topography includes various wavelengths with contributions from sub-lithosphere instabilities, plumes and other upper mantle heterogeneities potentially playing important roles in generating short-wavelength topography perturbations [Robinson and Parsons, 1988; King and Ritsema, 2000; Huang and Zhong, 2005; Faccenna and Becker, 2010; Yang and Leng, 2014; Schoonman et al., 2017]. However, our synthetic experiments suggest that the topographic recoverability is constrained by the present distribution of residual topography observations. Although short-wavelength topographic perturbations may be recovered locally with dense data coverages, increasing the maximum spherical degree from about degree 5 to degree 30 does not help the recovery of shorter-wavelength

features globally. On the other hand, the recovered residual topography power spectrum is generally consistent with that of dynamic topography for degrees less than 15. But the uncertainty is large and when the short-wavelength topographic relief is small while the measurement error is large, the spectral magnitude at short wavelengths may be strongly over estimated.

The choice of damping factors is important in parameter estimates, especially when the number of parameters inverted is large and the problem is underdetermined [Hartzell and Heaton, 1983; Rawlinson and Sambridge, 2003; Tarantola, 2005]. Both the error of the fit to observations and global synthetic-recovered topography differences depend on the choice of damping factors critically when the maximum expansion degree is larger than about 5 in the synthetic models investigated. The appropriate damping factors have a decreasing trend with increasing maximum expansion degree, suggesting that if one wants to fit the observation with more parameters, one should not smooth the model excessively. The small-amplitude long-wavelength residual topography derived by [Hoggard *et al.*, 2016; Hoggard *et al.*, 2017] is due to the large damping factor used. Although the conclusion that the residual topography amplitude reaches  $\sim 1\text{km}$  should not be influenced by a slight change of the damping factor, more objective ways may be needed in the future to yield the most appropriate maximum expansion degrees and damping factors automatically [Rawlinson and Sambridge, 2003]. The influence of truncating spherical harmonic expansions to a certain degree on long-wavelength residual topography recovery may also need to be considered [Trampert and Snieder, 1996].

## Acknowledgement

TY benefitted from the discussion with Judith Sippel on residual topography. The authors thank Malcolm Sambridge and two anonymous reviewers for reading the original manuscript and providing insightful suggestions. MG has been supported by the National Science Foundation through EAR-1358646, EAR-1600956, and EAR-1645775 and by Statoil ASA. LM and RDM were supported by Australian Research Council grants DP130101946 and IH130200012. Dynamic topography and the recovered long-wavelength residual topography data are listed in the supplements.

## References

- Čadež, O., and L. Fleitout (2003), Effect of lateral viscosity variations in the top 300 km on the geoid and dynamic topography, *Geophysical Journal International*, 152(3), 566-580.
- Cochran, J. R., and M. Talwani (1977), Free-air gravity anomalies in the world's oceans and their relationship to residual elevation, *Geophysical Journal International*, 50(3), 495-552.
- Colin, P., and L. Fleitout (1990), Topography of the ocean floor: Thermal evolution of the lithosphere and interaction of deep mantle heterogeneities with the lithosphere, *Geophysical Research Letters*, 17(11), 1961-1964.
- Conrad, C. P., and L. Husson (2009), Influence of dynamic topography on sea level and its rate of change, *Lithosphere*, 1(2), 110-120.
- Crough, S. T. (1978), Thermal origin of mid-plate hot-spot swells, *Geophysical Journal International*, 55(2), 451-469.
- Czarnota, K., M. Hoggard, N. White, and J. Winterbourne (2013), Spatial and temporal patterns of Cenozoic dynamic topography around Australia, *Geochemistry, Geophysics, Geosystems*, 14(3), 634-658.
- Davies, G. F., and F. Pribac (1993), Mesozoic seafloor subsidence and the Darwin Rise, past and present, in *The Mesozoic Pacific: Geology, Tectonics, and Volcanism*, edited by M. S. Pringle, W. W. Sager, W. V. Sliter and S. Stein, pp. 39-52, AGU, doi:10.1029/GM077p0039.
- Deschamps, F., R. Snieder, and J. Trampert (2001), The relative density-to-shear velocity scaling in the uppermost mantle, *Physics of the Earth and Planetary Interiors*, 124(3), 193-212.
- Faccenna, C., and T. W. Becker (2010), Shaping mobile belts by small-scale convection, *Nature*, 465(7298), 602-605.



- Flament, N., M. Gurnis, and R. D. Müller (2013), A review of observations and models of dynamic topography, *Lithosphere*, 5(2), 189-210.
- Gurnis, M. (1993), Phanerozoic marine inundation of continents driven by dynamic topography above subducting slabs, *Nature*, 364, 589-593.
- Gurnis, M., J. X. Mitrovica, J. Ritsema, and H. J. van Heijst (2000), Constraining mantle density structure using geological evidence of surface uplift rates: The case of the African superplume, *Geochemistry, Geophysics, Geosystems*, 1, doi:10.1029/1999GC000035.
- Hager, B., and M. Richards (1989), Long-wavelength variations in Earth's geoid: physical models and dynamical implications, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 328(1599), 309-327.
- Hager, B. H., R. W. Clayton, M. A. Richards, R. P. Comer, and A. M. Dziewonski (1984), Lower mantle heterogeneity, dynamic topography and the geoid, 313, 541-545.
- Hartzell, S. H., and T. H. Heaton (1983), Inversion of strong ground motion and teleseismic waveform data for the fault rupture history of the 1979 Imperial Valley, California, earthquake, *Bulletin of the Seismological Society of America*, 73(6A), 1553-1583.
- Hoggard, M., N. White, and D. Al-Attar (2016), Global dynamic topography observations reveal limited influence of large-scale mantle flow, *Nature Geoscience*, 9, 456-463, doi:10.1038/ngeo2709.
- Hoggard, M. J., J. Winterbourne, K. Czarnta, and N. White (2017), Oceanic Residual Depth Measurements, the Plate Cooling Model and Global Dynamic Topography, *Journal of Geophysical Research: Solid Earth*, doi: 10.1002/2016JB013457.
- Huang, J., and S. Zhong (2005), Sublithospheric small - scale convection and its implications for the residual topography at old ocean basins and the plate model, *Journal of Geophysical Research: Solid Earth*, 110(B5).
- Kaban, M., P. Schwintzer, and C. Reigber (2004), A new isostatic model of the lithosphere and gravity field, *Journal of Geodesy*, 78(6), 368-385.
- King, S. D., and J. Ritsema (2000), African hot spot volcanism: small-scale convection in the upper mantle beneath cratons, *Science*, 290(5494), 1137-1140.
- Le Stunff, Y., and Y. Ricard (1995), Topography and geoid due to lithospheric mass anomalies, *Geophysical Journal International*, 122(3), 982-990.
- Lithgow-Bertelloni, C., and M. Gurnis (1997), Cenozoic subsidence and uplift of continents from time-varying dynamic topography, *Geology*, 25(8), 735-738.
- Lithgow - Bertelloni, C., and M. A. Richards (1998), The dynamics of Cenozoic and Mesozoic plate motions, *Reviews of Geophysics*, 36(1), 27-78.
- Moucha, R., A. M. Forte, J. X. Mitrovica, D. B. Rowley, S. Quéré, N. A. Simmons, and S. P. Grand (2008), Dynamic topography and long-term sea-level variations: There is no such thing as a stable continental platform, *Earth and Planetary Science Letters*, 271(1), 101-108.
- Panasjuk, S. V., and B. H. Hager (2000a), Inversion for mantle viscosity profiles constrained by dynamic topography and the geoid, and their estimated errors, *Geophysical Journal International*, 143(3), 821-836.
- Panasjuk, S. V., and B. H. Hager (2000b), Models of isostatic and dynamic topography, geoid anomalies, and their uncertainties, *Journal of Geophysical Research: Solid Earth*, 105(B12), 28199-28209.
- Parsons, B., and S. Daly (1983), The relationship between surface topography, gravity anomalies, and temperature structure of convection, *Journal of Geophysical Research: Solid Earth*, 88(B2), 1129-1144.
- Pekeris, C. L. (1935), Thermal convection in the interior of the Earth, *Geophysical Journal International*, 3(8), 343-367, doi:10.1111/j.1365-246X.1935.tb01742.x.
- Rawlinson, N., and M. Sambridge (2003), Seismic traveltime tomography of the crust and lithosphere, *advances*

in *geophysics*, 46, 81-198.

Ricard, Y., M. Richards, C. Lithgow - Bertelloni, and Y. Le Stunff (1993), A geodynamic model of mantle density heterogeneity, *Journal of Geophysical Research: Solid Earth*, 98(B12), 21895-21909.

Robinson, E., and B. Parsons (1988), Effect of a shallow low - viscosity zone on the formation of midplate swells, *Journal of Geophysical Research: Solid Earth*, 93(B4), 3144-3156.

Rodríguez Tribaldos, V., N. J. White, G. Roberts, and M. J. Hoggard (2017), Spatial and temporal uplift history of South America from calibrated drainage analysis, *Geochemistry, Geophysics, Geosystems*.

Schoonman, C., N. White, and D. Pritchard (2017), Radial viscous fingering of hot asthenosphere within the Icelandic plume beneath the North Atlantic Ocean, *Earth and Planetary Science Letters*, 468, 51-61.

Schroeder, W. (1984), The empirical age - depth relation and depth anomalies in the Pacific Ocean Basin, *Journal of Geophysical Research: Solid Earth*, 89(B12), 9873-9883.

Shephard, G., L. Liu, R. Müller, and M. Gurnis (2012), Dynamic topography and anomalously negative residual depth of the Argentine Basin, *Gondwana Research*, 22(2), 658-663.

Spasojevic, S., and M. Gurnis (2012), Sea level and vertical motion of continents from dynamic earth models since the Late Cretaceous, *AAPG bulletin*, 96(11), 2037-2064.

Steinberger, B. (2007), Effects of latent heat release at phase boundaries on flow in the Earth's mantle, phase boundary topography and dynamic topography at the Earth's surface, *Physics of the Earth and Planetary Interiors*, 164(1), 2-20.

Tarantola, A. (2005), *Inverse problem theory and methods for model parameter estimation*, Society for Industrial and Applied Mathematics.

Trampert, J., and R. Snieder (1996), Model Estimations Biased by Truncated Expansions: Possible Artifacts in Seismic Tomography, *Science*, 271(5253), 1257-1260.

Wheeler, P., and N. White (2000), Quest for dynamic topography: Observations from Southeast Asia, *Geology*, 28(11), 963-966.

Winterbourne, J., N. White, and A. Crosby (2014), Accurate measurements of residual topography from the oceanic realm, *Tectonics*, 33(6), 982-1015.

Yang, T., and M. Gurnis (2016), Dynamic topography, gravity and the role of lateral viscosity variations from inversion of global mantle flow, *Geophysical Journal International*, 207(2), 1186-1202.


Yang, T., M. Gurnis, and S. Zahirovic (2016), Slab avalanche-induced tectonics in self-consistent dynamic models, *Tectonophysics*, <http://dx.doi.org/10.1016/j.tecto.2016.1012.1007>.

Yang, T., and W. Leng (2014), Dynamics of hidden hotspot tracks beneath the continental lithosphere, *Earth and Planetary Science Letters*, 401, 294-300.


Yue, H., T. Lay, J. T. Freymueller, K. Ding, L. Rivera, N. A. Ruppert, and K. D. Koper (2013), Supershear rupture of the 5 January 2013 Craig, Alaska (Mw 7.5) earthquake, *Journal of Geophysical Research: Solid Earth*, 118(11), 5903-5919.

Zahirovic, S., N. Flament, R. Dietmar Müller, M. Seton, and M. Gurnis (2016), Large fluctuations of shallow seas in low - lying Southeast Asia driven by mantle flow, *Geochemistry, Geophysics, Geosystems*, 17(9), 3589-3607.

Zhang, N., S. Zhong, and R. M. Flowers (2012), Predicting and testing continental vertical motion histories since the Paleozoic, *Earth and Planetary Science Letters*, 317, 426-435.



**Fig. 1** Synthetic topography field (a) represented by spherical harmonics up to degree 40 and its long-wavelength (degree 1-3) components (b). Green points represent the 2297 point-wise observations whose values are assumed known. Locations of these observations follow Hoggard et al. [2017]. The following quantities are shown as a function of maximum expanded spherical degrees and damping factors: (c) Error of the fit to observations; (d) average difference between recovered and synthetic topographies across the globe; (e) average difference between recovered and synthetic long-wavelength topographies across the globe; and (f) variance difference. Black dots in (c-f) correspond to the appropriate damping factors chosen from the misfit-roughness relationships in Fig. 2(a, b, c). Black dots in subsequent figures follow the same convention.



**Fig. 2** (a) Damping factor and recovered long-wavelength synthetic topography based on the misfit-roughness relationship with the maximum expansion degree fixed as 5. Left: Roughness of the recovered topography as a function of damping factor. Middle: error of the fit to observations (black) and variance difference (red) as a function of roughness. Right:

The recovered long-wavelength (degree 1-3) topography. The damping factor is chosen from the elbow of the misfit-roughness plot (black curve in the middle panel). (b, c) Same as (a) but the maximum expansion degree is fixed as 15 (b) and 30 (c), respectively. (d) Power spectrum as a function of spherical degree with the damping factor fixed at 0.84 (Fig. 1a). Different colors represent spherical expansions with different maximum degrees, green represents the synthetic. (e) Recovered long-wavelength topography with the maximum expansion degree as 30 and damping factor as 0.84.







Fig. 3

**Fig. 3** Long-wavelength topography recovery for synthetic models with different power spectra. (a-e) Spectral power does not decrease with increasing degree (exponent  $p=0.0$ ). (f-j) Spectral power decreases faster with increasing degrees as compared with the case in Fig. 1

(exponent  $p=-0.8$ ). (a, f) synthetic topography, (b, g) their long-wavelength components, (c, h) error of the fit to observations and variance difference, (d, i) damping factor from misfit-roughness plot, (e, j) the recovered long-wavelength topographies.



**Fig. 4** Comparisons between mantle flow model predicted dynamic topography and recent estimated residual topography in the oceanic realm. (a) Dynamic topography and (b) its long-wavelength (degree 1-3) components are from *Yang and Gurnis [2016]*. (c) Error of the fit to 2297 residual topography observations. (d-f) Inferred long-wavelength residual topography based on 2297 observations with the maximum expansion degree as 5, 12 and 30. The corresponding damping factor for each degree is chosen from the elbow of the misfit-roughness relationships (Fig. S5). In (a, b), circles represent the 1161 observations which have both precise sedimentary and crustal corrections, while upward- and downward-pointing triangles represent the remaining points for which residual topography might be over- or under- estimated, respectively [*Hoggard et al., 2017*].